Searches for stochastic gravitational-wave backgrounds

Lecture 1 Les Houches Summer School 19 July 2018



Joe Romano University of Texas Rio Grande Valley **Texas Tech University**





Connection to other lectures

- Astrophysical sources I. Mandel (last week)
- Cosmological sources V. Mandic (starting tomorrow)
- Data analysis A. Weinstein (last week), J. Veitch (this week)
- Pulsar timing N. Cornish (next week)
- LISA detector / science M. Hewitson, A. Sesana (next week)

Some references (not complete)



- M. Maggiore "Gravitational-wave experiments and early universe cosmology" (2000)
- C. Caprini, D. Figueroa "Cosmological backgrounds of gravitational waves" (2018)
- T. Regimbau "The astrophysical stochastic gravitational-wave backgrounds" (2011)
- J. Romano, N. Cornish "Detection methods for stochastic gravitational-wave backgrounds: a unified treatment" (2017)
- R. Smith, E. Thrane "Optimal search for an astrophysical gravitational-wave background" (2018)
- Plus recent observational papers from LIGO, Virgo, pulsar timing arrays, etc., quoting upper limits on the strength of stochastic gravitational-wave backgrounds

B. Allen - "The stochastic gravitational-wave background: sources and detection," from Les Houches School in

Resources

https://github.com/josephromano/leshouches

Slides

• Exercises (suggested)

Solutions

Code examples (ipython notebooks)

Plan for lectures

Today: Overview / Basics

- 1. Motivation / context
- 2. Different types of stochastic backgrounds
- 3. Characterizing a stochastic GW background
- 4. Correlation methods
- 5. Some simple examples

Tomorrow: Details / Example

- 1. Non-trivial detector response
- 2. Non-trivial correlated response
- 3. What to do in the absence of correlations (e.g., for LISA)?
- 4. Frequentist and Bayesian methods
- 5. Example: searching for the background from BBH mergers

1. Motivation

Ultimate goal: produce GW analogue of CMB sky map





But there's a long road ahead...

1965: Penzias & Wilson

Measurements of the effective zenith noise temperature of the 20-foot horn-reflector antenna (Crawford, Hogg, and Hunt 1961) at the Crawford Hill Laboratory, Holmdel, New Jersey, at 4080 Mc/s have yielded a value about 3.5° K higher than expected. The excess temperature is, within the limits of our observations, isotropic, unpolarized, an free from seasonal variations (July, 1964–April, 1965). A possible explanation for th observed excess noise temperature is the one given by Dicke, Peebles, Roll, and Wilkinso (1965) in a companion letter in this issue.



2013: WMAP, Planck

1992: COBE

2018: And we haven't detected the isotropic component of the GW background yet!

A MEASUREMENT OF EXCESS ANTENNA TEMPERATURE AT 4080 Mc/s (4080 Mc/s <-> 7.35 cm)



At least we've detected other GW signals...

- 5 (or 6) binary black hole (BBH) mergers and binary neutron star (BNS) merger
- with similar detections expected in O3, ...
- very strong events!!



Observation of Gravitational Waves from a Binary Black Hole Merger

B. P. Abbott *et al.**

(LIGO Scientific Collaboration and Virgo Collaboration) (Received 21 January 2016; published 11 February 2016)

On September 14, 2015 at 09:50:45 UTC the two detectors of the Laser Interferometer Gravitational-Wave Observatory simultaneously observed a transient gravitational-wave signal. The signal sweeps upwards in frequency from 35 to 250 Hz with a peak gravitational-wave strain of 1.0×10^{-21} . It matches the waveform predicted by general relativity for the inspiral and merger of a pair of black holes and the ringdown of the resulting single black hole. The signal was observed with a matched-filter signal-to-noise ratio of 24 and a false alarm rate estimated to be less than 1 event per 203 000 years, equivalent to a significance greater than 5.1 σ . The source lies at a luminosity distance of 410^{+160}_{-180} Mpc corresponding to a redshift $z = 0.09^{+0.03}_{-0.04}$. In the source frame, the initial black hole masses are $36^{+5}_{-4}M_{\odot}$ and $29^{+4}_{-4}M_{\odot}$, and the final black hole mass is $62_{-4}^{+4}M_{\odot}$, with $3.0_{-0.5}^{+0.5}M_{\odot}c^2$ radiated in gravitational waves. All uncertainties define 90% credible intervals. These observations demonstrate the existence of binary stellar-mass black hole systems. This is the first direct detection of gravitational waves and the first observation of a binary black hole merger.







...and we expect many more weaker signals

• individually undetectable (subthreshold)

- but detectable as a collectivity via their common influence on multiple detectors
- combined signal described statistically stochastic gravitational-wave background

Exercise 1: Verify that the expected total rate of stellar-mass BBH mergers is between ~1 per minute and a few per hour.



Potentially detectable with advanced LIGO/Virgo



(PRL 120, 091101, 2018)

3)		
	-		
•	•		
		-	
			-
			_
1			
			-
			1
			1
			-
			-
			1
1			

Bigger picture of GWs — sources & detectors





2. Different types of stochastic GW backgrounds

(i) Stochastic backgrounds can differ in spatial distribution

(statistically) isotropic



(like cosmic microwave background)

<u>anisotropic</u>





(ii) They can also differ in temporal distribution and amplitude









Example: Rate estimates and signal durations imply BNS "confusion" & BBH "popcorn" for LIGO / Virgo





Combined BBH / BNS background signal







3. Characterizing a stochastic GW background

Definition of a stochastic background

- Superposition of signals too weak or too numerous to individually detect
- Looks like noise in a single detector
- Characterized statistically in terms of moments (ensemble averages) of the metric perturbations





Plane wave expansion of metric perturbations

$$h_{ab}(t, \overrightarrow{x}) = \int_{-\infty}^{\infty} \mathrm{d}f \int \mathrm{d}^2 \Omega_{\hat{n}} \sum_{A=+,\times} h_A(f, \hat{n}) e_{ab}^A(t, dx)$$

Polarization tensors:





Quadratic expectation values specify different types of Gaussian stochastic backgrounds

<u>Unpolarized, stationary</u> <u>isotropic:</u>

 $\left\langle h_A(f,\hat{n})h_{A'}^*(f',\hat{n}')\right\rangle = -$

<u>Unpolarized, stationary,</u> <u>anisotropic:</u> $\left\langle h_A(f,\hat{n})h_{A'}^*(f',\hat{n}')\right\rangle = -\frac{1}{2}$

where

 $S_h(f) =$

power spectral density (Hz-1)

$$S_h(f) = \frac{3H_0^2}{2\pi^2} \frac{\Omega_{\rm gw}(f)}{f^3}$$

Exercise 2: Derive the above relationship.

energy density spectrum (dimensionless)

$$\Omega_{\rm gw}(f) \equiv -\frac{1}{\rho}$$

$$o_c \equiv \frac{3H_0^2 c^2}{8\pi G}$$

$$\frac{1}{16\pi} S_h(f) \delta(f - f') \delta_{AA'} \delta^2(\hat{n}, \hat{n}')$$

$$\frac{1}{4} \mathcal{P}(f, \hat{n}) \delta(f - f') \delta_{AA'} \delta^{2}(\hat{n}, \hat{n}')$$

$$\int d^{2}\Omega_{\hat{n}} \mathcal{P}(f, \hat{n})$$

characteristic strain (dimensionless)

$$\frac{\mathrm{d}\rho_{\mathrm{gw}}}{\mathrm{d}\ln f} = \frac{f}{\rho_c} \frac{\mathrm{d}\rho_{\mathrm{gw}}}{\mathrm{d}f} \qquad \qquad h_c(f) \equiv \sqrt{fS_h(f)} = A_\alpha \left(\frac{f}{f_{\mathrm{ref}}}\right)$$

$$\rho_{gw} = \frac{c^2}{32\pi G} \langle \dot{h}_{ab}(t, \vec{x}) \dot{h}^{ab}(t, \vec{x}) \rangle$$



"Phinney formula": Calculating $\Omega_{gw}(f)$ for an astrophysical background

Recall:

$$\Omega_{gw}(f) \equiv \frac{1}{\rho_c} \frac{d\rho_{gw}}{d\ln f} = \frac{f}{\rho_c} \frac{d\rho_{gw}}{df}$$

For a collection of sources:

$$\Omega_{gw}(f) = \frac{1}{\rho_c} \int_0^\infty dz \, n(z) \frac{1}{1+z} \left(f_s \frac{dE_{gw}}{df_s} \right) \Big|_{f_s = f(1+z)}$$

$$f_s = f(1+z)$$
In terms of event rate:

$$n(z) \, dz = R(z) \left| \, dt \right|_{\rm s}$$

 $\left| \frac{\mathrm{d}t}{\mathrm{d}z} \right|_{s} = \frac{1}{(1+z)^{s}}$

Exercise 3: Verify the above expression for |dt/dz| as well as the "Phinney formula" in terms of the rate R(z).

$$\Omega_{gw}(f) = \frac{f}{\rho_c H_0} \int_0^\infty dz \, R(z) \frac{1}{(1+z)E(z)} \left(\frac{dE_{gw}}{df_s}\right) \bigg|_{f_s = f(1+z)}$$

astro-ph/0108028

A practical theorem on gravitational wave backgrounds

E.S. Phinney^{*}

Theoretical Astrophysics, 130-33 Caltech, Pasadena, CA 91125, USA

ABSTRACT

There is an extremely simple relationship between the spectrum of the gravitational wave background produced by a cosmological distribution of discrete gravitational wave sources, the total time-integrated energy spectrum of an individual source, and the present-day comoving number density of remnants. Stated in this way, the background is entirely independent of the cosmology, and only weakly dependent on the evolutionary history of the sources. This relationship allows one easily to compute the amplitude and spectrum of cosmic gravitational wave backgrounds from a broad range of astrophysical sources, and to evaluate the uncertainties therein.

$$\frac{1}{z)H_0E(z)} \qquad E(z) = \sqrt{\Omega_m(1+z)^3 + \Omega_\Lambda} \quad \longleftarrow \quad \text{cosmology}$$



Simple example: circular binaries

Units: G = c = 1Kepler's law: $\omega^2 r^3 = GM \implies r \sim M^{1/3} \omega^{-2/3}, \quad \dot{r} \sim -r \dot{\omega}/\omega$ **Energy balance:** $\frac{dE_{gw}}{dt} = -\frac{dE_{orb}}{dt}$ $\implies \frac{dE_{gw}}{dt} \sim -M\mu \dot{r}/r^2 \sim \mathcal{M}_c^{5/3} \omega^{-1/3} \dot{\omega}$ $\implies \frac{dE_{gw}}{df} = \frac{dt}{df} \frac{dE_{gw}}{dt} \sim \frac{1}{\dot{\omega}} \frac{dE_{gw}}{dt} \sim \mathcal{M}_c^{5/3} f^{-1/3}$ $\Omega_{\rm gw}(f) \propto f^{2/3}, \quad h_c(f) \propto f^{-2/3}$







$$\Omega_{\rm gw}(f) \propto f^{2/3}, \quad h_c(f) \propto f^{-2/3}$$

4. Correlation methods

Single vs. multiple detectors

- Initial discovery of the CMB was in a single detector (excess noise that could not be attributed to any known noise source)
- Ground-based detectors currently aren't sensitive enough for expected GW backgrounds to stand out above instrumental noise (LISA is another story!)
- Instead, look for evidence of a common disturbance in multiple detectors -> crosscorrelation
- Signal might be weak, but you can build up SNR by correlating for long periods of time

A MEASUREMENT OF EXCESS ANTENNA TEMPERATURE AT 4080 Mc/s

Measurements of the effective zenith noise temperature of the 20-foot horn-reflector antenna (Crawford, Hogg, and Hunt 1961) at the Crawford Hill Laboratory, Holmdel, New Jersey, at 4080 Mc/s have yielded a value about 3.5° K higher than expected. This excess temperature is, within the limits of our observations, isotropic, unpolarized, and free from seasonal variations (July, 1964-April, 1965). A possible explanation for the observed excess noise temperature is the one given by Dicke, Peebles, Roll, and Wilkinson (1965) in a companion letter in this issue.

(Penzias & Wilson, 1965)





Cross-correlation: basic idea

Data from two detectors:



common GW signal component

Expected value of cross-correlation:

 $\langle \hat{C}_{12} \rangle = \langle d_1 d_2 \rangle = \langle h^2 \rangle + \langle n_1 d_2 \rangle$

Assuming detector noise is uncorrelated:



$$= h + n_1$$
$$= h + n_2$$

$$|n_2\rangle + \langle hn_2\rangle + \langle n_1h\rangle = \langle h^2\rangle + \langle n_1n_2\rangle$$

$$= \langle h^2 \rangle \equiv S_h$$

$$\hat{S}_h \equiv \hat{C}_{12} = \frac{1}{N} \sum_{i=1}^N d_{1i} d_{2i}$$
 $d_{1i} = h_i + n_{1i}, \quad d_{2i} = h_i + n_{2i}$

Expected value:

$$\begin{split} \mu &\equiv \langle \hat{C}_{12} \rangle = \frac{1}{N} \sum_{i=1}^{N} \langle d_{1i} d_{2i} \rangle = \frac{1}{N} \sum_{i=1}^{N} \langle h_i^2 \rangle = S_h \\ \text{Variance:} \qquad \sigma^2 &\equiv \langle \hat{C}_{12}^2 \rangle - \langle \hat{C}_{12} \rangle^2 = \left(\frac{1}{N}\right)^2 \sum_{i=1}^{N} \sum_{j=1}^{N} \left(\langle d_{1i} d_{2i} d_{1j} d_{2j} \rangle - \langle d_{1i} d_{2i} \rangle \langle d_{1j} d_{2j} \rangle \right) \\ &\downarrow \langle abcd \rangle = \langle ab \rangle \langle cd \rangle + \langle ac \rangle \langle bd \rangle + \langle ad \rangle \langle bc \rangle \\ &= \left(\frac{1}{N}\right)^2 \sum_{i=1}^{N} \sum_{j=1}^{N} \left(\langle d_{1i} d_{1j} \rangle \langle d_{2i} d_{2j} \rangle + \langle d_{1i} d_{2j} \rangle \langle d_{2i} d_{1j} \rangle \right) \\ &= \left(\frac{1}{N}\right)^2 \sum_{i=1}^{N} \sum_{j=1}^{N} \left(S_1 \delta_{ij} S_2 \delta_{ij} + S_h \delta_{ij} S_h \delta_{ij} \right) \qquad \text{where} \qquad \frac{S_1}{S_2} \\ &= \left(\frac{1}{N}\right)^2 \sum_{i=1}^{N} \left(S_1 S_2 + S_h^2 \right) = \frac{1}{N} \left(S_1 S_2 + S_h^2 \right) \simeq \frac{1}{N} S_1 S_2 \\ \text{SNR:} \qquad \rho \equiv \frac{\mu}{\sigma} \simeq \frac{S_h}{\sqrt{S_1 S_2 / N}} \qquad \Longrightarrow \qquad \left[\rho \simeq \sqrt{N} \frac{S_h}{S_n} \right] \qquad \text{where} \qquad \sqrt{S_1 S_2} \approx \sqrt{S_{n_1} S_{n_2}} \equiv S_n \end{split}$$

Worked example: N samples, white GWB in white detector noise

(common GW signal, uncorrelated detector noise)

$$= S_h$$

(8282482)	where	$S_1 = S_{n_1} + S_h$
$ij S_2 O_{ij} + S_h O_{ij} S_h O_{ij}$		$S_2 = S_{n_2} + S_h$

Cross-correlation estimators / optimal filtering

More generally:

$$\hat{S}_{h} = \int_{-T/2}^{T/2} \mathrm{d}t \int_{-T/2}^{T/2} \mathrm{d}t' \, dt' \, dt$$

Choose Q to maximize SNR for fixed spectral shape:

correlation coeff (overlap) between two detectors

$$\langle \tilde{h}_1(f)\tilde{h}_2^*(f')\rangle = \frac{1}{2}\delta(f-f')\Gamma_{12}(f)S_h(f)$$

Exercise 4: Verify the expression for the optimal filter function Q(f).

 $d_1(t)d_2(t')Q(t,t') \qquad \text{if stationary} \\ Q(t,t') = Q(t-t')$

 $_{T}(f-f')\tilde{d}_{1}(f)\tilde{d}_{2}^{*}(f')\tilde{Q}^{*}(f')$



30

5. Some simple examples

(code to simulate data and do the analyses are on github repository)

(i) White GWB in white detector noise



expected and estimated values of power agree to 3.5%, within 1 sigma



optimally filtered CC SNR = 2.9

(ii) Confusion-limited BNS GWB in white detector noise



expected and estimated values of power agree to 2.7%, within 1 sigma

optimally filtered CC SNR = 12





(iii) Two-component GWB in white detector noise (GWB = white GWB + confusion-limited BNS GWB)



optimal-filtering for each component separately: white GWB: 48% overestimate, > 1 sigma BNS GWB: 6.9% overestimate, within 1 sigma

separate analyses overestimate strength of each GWB component and underestimate error bars



joint multi-component analysis: white GWB: agreement to 7.3%, SNR=1.4 BNS GWB: agreement to 3.8%, SNR=6.0

joint analysis properly takes into account the covariance between the component spectral shapes

Parida et al, JCAP 024, 2106

Joint multi-component analysis

	$F_{\alpha\beta} = \int_{-\infty} \mathrm{d}f$
Exercise 5 : Verify the expression for the ML estimators.	$F \equiv M$
Maximum-likelihood estimators:	$\hat{A} = F$
Likelihood function:	$p(\hat{C} A$
Noise covariance matrix:	N ₁₂ (f, f'
Expectation value:	$\langle \hat{C}_{12}(f)$
Data are cross-correlation estimates:	$\hat{C}_{12}(f)$



$$\begin{aligned} \hat{C}' \end{pmatrix} &\equiv \langle \hat{C}_{12}(f) \hat{C}_{12}^*(f') \rangle - \langle \hat{C}_{12}(f) \rangle \langle \hat{C}_{12}^*(f') \rangle \\ &\simeq \delta_{ff'} P_1(f) P_2(f) \end{aligned}$$

$$(A, N) \propto \exp\left[-\frac{1}{2}(\hat{C} - MA)^{\dagger}N^{-1}(\hat{C} - MA)\right]$$

 $T^{-1}X$

 $M^{\dagger}N^{-1}M, \qquad X \equiv M^{\dagger}N^{-1}\hat{C}$

 $H_{\alpha}(f)\Gamma_{12}^{2}(f)H_{\beta}(f)$ $P_1(f)P_2(f)$

(noise-weighted inner product of spectral shapes; inverse covariance matrix for A)

Fisher matrix



end lecture 1



extra slides



What are the prospects for detection?

No detections yet; only upper limits on strength of background in different freq bands.



Friedmann-Robertson-Walker line element:

 $ds^{2} = -c^{2} dt^{2} + a^{2}(t) \left(dr^{2} + r^{2} d\theta^{2} + r^{2} \sin^{2} \theta d\phi^{2} \right)$

Redshifted frequencies and time-intervals:

$$1 + z = \frac{a(t_0)}{a(t_s)} \longrightarrow 1 + z = \frac{1}{a(t_s)}, \quad a(t_0) \equiv 1$$

$$f_s = (1 + z)f_0$$

$$\Delta t_0 = (1 + z)\Delta t_s$$

$$\frac{dt}{dz} = \frac{1}{(1 + z)H_0E(z)}$$

$$f_{riedmann equation}$$

$$\frac{\dot{d}}{dz} = H(t) = H_0E(z)$$

$$E(z) = \sqrt{\Omega_m(1 + z)^3 + \Omega_m}$$

$$f_{riedmann equation}$$

$$f_{riedmann equation}$$



